



Mathematics and CS Seminar

Waists of balls in different spaces

Arseniy Akopyan (IST Austria)

Host: Jan Maas

Gromov and Memarian (2003--2011) have established the waist inequality asserting that for any continuous map f from the sphere S^n to \mathbb{R}^{n-k} there exists a fiber $f^{-1}(y)$ such that every t -neighborhood has measure at least the measure of the t -neighborhood of an equatorial subsphere S^k of S^n . Going to the limit we may say that the $(n-k)$ -volume of the fiber $f^{-1}(y)$ is at least that of the standard sphere S^k . We extend this limit statement to the exact bounds for balls in spaces of constant curvature, tori, parallelepipeds, projective spaces and other metric spaces. By the volume of preimages for a non-regular map f we mean its lower Minkowski content, some new properties of which will be also presented in the talk. (based on the joint work with Roman Karasev and Alfredo Hubard)

Thursday, February 8, 2018 04:00pm - 06:00pm

IST Austria Campus Big Seminar room Ground floor / Office Bldg West (I21.EG.101)



This invitation is valid as a ticket for the IST Shuttle from and to Heiligenstadt Station. Please find a schedule of the IST Shuttle on our webpage: http://ist.ac.at/fileadmin/user_upload/pdfs/IST_shuttle_bus.pdf The IST Shuttle bus is marked IST Shuttle (#142) and has the Institute Logo printed on the side.