



FriSBI

Liquid plug formation in human airways

Francesco Romanò (University of Michigan)

Host: Gasper Tkacik, Edouard Hannezo

The airway of human lungs articulate from the windpipe in a network of bifurcating branches known as bronchi and bronchioles. At each airway generation, i.e. when a branch bifurcates, the airway cross section reduces, up to becoming microscopic when it connects to the alveoli. Considering that human lung airways are coated lined with a liquid made out of mucus and serous, after 8 or 9 generations the surface tension between this liquid layer and the air might induce a Rayleigh-Plateau instability of a thick-enough liquid film. This phenomenon, known as airway closure, creates a liquid plug which blocks the airway halting distal gas exchange. Consequence of the airway closure are the flow-induced high stress levels on the wall, which is the location of airway epithelial cells. Relevant conditions for human lungs are simulated, taking into account ordinary and pathological parameters. Our numerical prediction is able to capture the physical process from pre- to post-coalescence, whereas previous studies have been limited to pre-coalescence only. Therefore, we can study the effect of the topological change, and we discovered that during coalescence, a high level of stress and stress gradients is exerted on the epithelial cells. We find that wall stresses during the post-coalescence phase can be in the range of 300% to 600% greater than pre-coalescence values. Hence, airway closure qualifies as a cause of sub-lethal or lethal responses for the epithelial cells.

Friday, September 14, 2018 03:00pm - 04:00pm

IST Austria Campus Mondi Seminar Room 3, Central Building



This invitation is valid as a ticket for the IST Shuttle from and to Heiligenstadt Station. Please find a schedule of the IST Shuttle on our webpage: <https://ist.ac.at/en/campus/how-to-get-here/> The IST Shuttle bus is marked IST Shuttle (#142) and has the Institute Logo printed on the side.