



Physical Sciences Seminar

From Single to interacting Polarons in Bose-Einstein condensates

Luis Peña Ardila

Aarhus University

Host: Misha Lemeshko

Mobile impurities in a Bose-Einstein condensate can form quasi-particles termed Bose-Polarons. In this talk I show how these quasi-particles are originated when a single impurity is dressed by the excitations of the quantum bosonic bath. The moststriking advantage of these polarons is the huge degree of controllability of thecoupling strength between the impurity and the bosonic bath. Thus, one can realize polarons from weak all the way up to the strong interacting regime [1,2]. For strong interactions two polaron can bind together forming bound bipolarons states. They emerge due to the induced non-local interaction mediated by density oscillations of the bath [3,4]. Polarons and bipolarons in ultra-cold quantum gases could be used as a robust platform for quantum simulation in a condensed matter context.REFERENCES.[1] N. B. Jrgensen, L. Wacker, K. T. Skalmstang, M. M. Parish, J. Levinsen, R. S. Christensen, G. M. Bruun, and J. J. Arlt. Phys. Rev. Lett. 117 . 055302 (2016).[2] M-G Hu, M. J. V. de Graaff, D. Kedar, J. P. Corson, E. A. Cornell, and D. S. Jin. Phys. Rev. Lett. 117, 055301 (2016).[3] L. A. P. Ardila and S. Giorgini. Phys. Rev. A 92, (2015).[4] A. Camacho-Guardian, Luis. A. Pea Ardila, Thomas Pohl and G. M. Bruun. Phys. Rev. Lett. 121, 013401 (2018)

Tuesday, January 15, 2019 11:00am - 12:30pm

Big Seminar room Ground floor / Office Bldg West (I21.EG.101)



This invitation is valid as a ticket for the ISTA Shuttle from and to Heiligenstadt Station.

Please find a schedule of the ISTA Shuttle on our webpage:

<https://ista.ac.at/en/campus/how-to-get-here/> The ISTA Shuttle bus is marked ISTA Shuttle (#142) and has the Institute Logo printed on the side.