

Mathematics and CS Seminar

Local inhomogeneous circular law

Johannes Alt

IST Austria

Host: Laszlo Erdös

The density of eigenvalues of large random matrices typically converges to a deterministic limit as the dimension of the matrix tends to infinity. In the Hermitian case, the best known examples are the Wigner semicircle law for Wigner ensembles and the Marchenko-Pastur law for sample covariance matrices. In the non-Hermitian case, the most prominent result is Girko s circular law: The eigenvalue distribution of a matrix X with centered, independent entries converges to a limiting density supported on a disk. Although inhomogeneous in general, the density is uniform for identical variances. In this special case, the local circular law by Bourgade et. al. shows this convergence even locally on scales slightly above the typical eigenvalue spacing. In the general case, the density is obtained via solving a system of deterministic equations. In my talk, I explain how a detailed stability analysis of these equations yields the local inhomogeneous circular law in the bulk spectrum for a general variance profile of the entries of X. This result was obtained in joint work with László Erdös and Torben Krüger.

Thursday, March 9, 2017 04:00pm - 06:00pm

Seminar room Big Ground floor / Office Bldg West (I21.EG.101)



This invitation is valid as a ticket for the ISTA Shuttle from and to Heiligenstadt Station. Please find a schedule of the ISTA Shuttle on our webpage: https://ista.ac.at/en/campus/how-to-get-here/ The ISTA Shuttle bus is marked ISTA Shuttle (#142) and has the Institute Logo printed on the side.

www.ista.ac.at | Institute of Science and Technology Austria | Am Campus 1 | 3400 Klosterneuburg